

Rain on snow responses to climate warming in the Pyrenees

Josep Bonsoms ¹, Juan Ignacio López-Moreno ², Esteban Alonso-González ³

1. Department of Geography, Universitat de Barcelona, Barcelona, Spain.

2. Instituto Pirenaico de Ecología (IPE-CSIC), Campus de Aula Dei, Zaragoza, Spain.

3. Centre d'Etudes Spatiales de la Biosphère (CESBIO), Université de Toulouse, CNES/CNRS/IRD/UPS, Toulouse, France.

Introduction

A warmer atmosphere is expected to decrease the snowfall ratio, leading to rain on snow (ROS) events, increasing the snow runoff and triggering natural hazards, such as flash flood events.


ROS is a complex phenomena, since a warmer atmosphere reduces also the snow covered season and therefore the ROS occurrence. Previous studies has shown different ROS sensitivities due to warming, depending on the elevation and the month of the season (McCabe et al 2007, Beniston and Stoffel, 2016; Morán-Tejeda et al 2016; López-Moreno et al., 2021). However, in the Pyrenees, only Corripio and López-Moreno (2017) analyzed the impact of ROS in the Surface Energy Balance for a single ROS event.

In this work we aim to analyzed the ROS sensitivity due to climate change in the Pyrenees.

ROS sensitivity to climate warming is analyzed for low, mid and high elevation sectors of the mountain range.

ROS sensitivity is analyzed through three variables:

- 1 ROS Frequency
- 2 Rainfall intensity during ROS
- 3 Snow ablation during ROS



Atmospheric data

Système d'Analyse Fournissant des Renseignements Adaptés à la Nivologie (SAFRAN).

Flat slopes at low, mid and high elevation ranges and Pyrenean massifs (Figure 1) at hourly resolution (Vernay et al., 2021)

Snow model

Physical-based and mass balance snow model: Flexible Snow Model (FSM2; Essery, 2015).

FSM2 resolves the SEB simulating the state of the snowpack.

FSM2 configuration includes runoff, internal snow processes, refreeze and thermal conductivity, quantified as function of the snow density. Atmospheric stability is simulated as function of the Richardson number.

ROS detection and climate sensitivity

ROS detection : daily rainfall > 10 mm and snow depth >= 0.1 (López-Moreno et al., 2021)

Climate sensitivity (1980 – 2019 baseline climate) analyzed through a delta-change methodology:

Precipitation variation from -10 to 10 % and increments of temperature from +1 to +4°C, according to climate change projections (Amblar-Francés et al., 2020).

Geographical setting




Figure 1. Geographical area analyzed in this work.

Location

The Pyrenees is located in the South of Europe (42°N-43°N to 2°W-3°E), between the Atlantic Ocean (West) and the Mediterranean Sea (East).

Elevation increases towards the central area, where the highest peak is found (Aneto, 3.404 m asl).

Climatology

Transition belt between continental and subtropical climate.

Precipitation


High spatial variability, ~ 1000 mm/year. Maximum (~ 2000 mm/year) is found in the central mountain summits. Increases/decreases from West (North) to East (South)

Temperature

Adiabatic rate: ~ 0.55°C/100 m (Navarro-Serrano and López-Moreno, 2018).

Annual isotherm of 0°C ~ 2750 – 2950 m (López-Moreno and García-Ruiz, 2004).

Snow ablation Net radiation (55 %), latent (32 %) and sensible heat flux (13 %) (Bonsoms et al., 2022).



Credits: UB and CESBIO

Results



Figure 2. Sensitivity of ROS frequency and rainfall quantity




Figure 3. Snow depth and snowfall fraction evolution.

Rainfall quantity during ROS

For all elevation zones: general increase in Winter.

At low and mid elevation: decreases during Spring.

At high elevation: general increases specially during Summer and Spring.

Spatial differences

Western Pyrenees:

- > ROS per year
- > Average daily snow ablation
- > rainfall quantity during ROS events

ROS frequency

For all elevation aras: general increase in Winter.

At low and mid elevation: ROS frequency generally decreases (increases) during Autumn (Spring) for > +2°C.

At high elevation: large increases during Spring (western area). Decreases for +2°C during Summer.

Snow ablation

Higher average daily snow ablation during ROS events than during average snow ablation episodes.

Small variation in the average daily snow ablation due to climate warming.

No spatial differences found with respect to baseline climate.




Figure 4. Spatial distribution of the rainfall intensity during ROS events




Figure 5. ROS frequency spatial distribution

Table 1. Increase of the rainfall intensity during ROS events

Elevation	Season	Baseline	+1°C	+2°C	+3°C	+4°C
Low	Autumn	14.6	+8.5	+11.6	+12.3	+11.1
	Winter	17.2	+3.9	+7	+8.3	+8.4
	Spring	16.3	+4.4	+6.7	+7.8	+7.7
Mid	Autumn	No ROS	No ROS	No ROS	No ROS	No ROS
	Winter	17	+7.4	+11	+8.4	+9
	Spring	16.7	+5.1	+6.8	+8.8	+10.6
High	Autumn	15.7	+4.6	+7.2	+8.8	+8.8
	Summer	10.1	+10.5	+14.7	+8.8	+10.6
	Spring	14.8	+8.4	+13.6	+16.1	+16
High	Autumn	10.4	+9.2	+14	+16	+16.7
	Winter	10.7	+6.4	+10.6	+12.8	+15.3
	Summer	25.2	-1.4	-0.1	+3.3	+16




Figure 6. Average daily snow ablation during (a) all days and (b) ROS days.

ROS frequency is key controlled by snow depth duration




Figure 7. Pearson's correlation between ROS frequency (days/year) and yearly (a) HS duration, (b) Snowfall fraction and (c) temperature.

Conclusions

At low elevation, ROS events will decrease during Autumn and Spring.

ROS events and rainfall quantity during ROS increase in a warmer climate at high elevations.

High average daily snow ablation during ROS events.

Seasonal snow depth duration controls yearly ROS frequency

Future research

ROS interannual variability and sensitivity between extreme snow seasons

ROS physical process

Extreme events evaluation

Short term hydrological response

References

Amblar-Francés, M.P., Ramos-Cabado, P., Sánchez-Liada, J., Hernández-Lázaro, A., Peral, García, M.C., Navasacs, B., Domínguez-Alonso, M., and Rodríguez-Camino, E.: High resolution climate change projections for the Pyrenees region. *Adv. Sci. Res.*, 17, 191–208, 2020.

Beniston, M., Stoffel, M.: Rain-on-snow events, floods and climate change in the Alps: events may increase with warming up to 4 °C and decrease thereafter. *Science of the Total Environment*, 571:228–236, 2016.

Bonsoms, J., López-Moreno, J.I., González, S., Oliva, M.: Increase of the energy available for snow ablation and its relation with atmospheric circulation. *Atmospheric Research*, 275:10628, 2022.

Compte, J. G., and López-Moreno, J. I.: Analysis and predictability of the hydrological response of mountain catchments to heavy rain on snow events: A case study in the Spanish Pyrenees. *Hydrology*, 4(2), 20, 2017.

Essery, R.: A factorial snowpack model (FSM 1.0). *Geoscientific Model Development*, 8(12), 3667–3676, 2015.

López-Moreno, J.I., García-Ruiz, J.M.: Influence of snow accumulation and snowmelt on streamflow in the Central Spanish Pyrenees. *Hydrological Sciences Journal*, 49 (5): 787–802, 2004.

López-Moreno, J.I., Pomroy, J.W., Morán-Tejeda, E., Revuelto, J., Navarro-Serrano, F.M., Vidaller, I., Alonso-González, E.: Changes in the frequency of global high mountain rain-on-snow events due to climate warming. *Environ. Res. Lett.*, 16, 044021, 2021.

McCabe, G. J., Clark, M. P., Hay, L. E.: Rain-on-snow events in the western United States. *Int. J. Meteorol. Soc.* 35, 319–328, 2007.

Morán-Tejeda, E., López-Moreno, J.I., Stoffel, M., Beniston, M.: Rain-on-snow events in Switzerland: recent observations and projections for the 21st century. *Clim Res* 71: 111–125, 2016.

Navarro-Serrano, F. and López-Moreno, J.I.: Spatio-temporal analysis of snowfall events in the Spanish Pyrenees and their relationship to atmospheric circulation. *Cuad. Invest. Geogr.*, 43 (1), 233–264, 2017.

Vernay, M., Lafaysse, M., Morin, D., Hagenmüller, P., Nival, R., Samadizadeh, R., Verfaillie, D., and Morin, S.: The S2M meteorological and snow cover reanalysis over the French mountainous areas, description and evaluation (1958–2020). *Earth Syst. Sci. Data Discuss*, 2021.

